

Macrolides in practice for swine



02-2026

Active component	Tylosin	Tylvalosin	Tilmicosin
	Natural macrolide, produced by <i>Streptomyces fradiae</i>	Semi-synthetic derivate of tylosin	Semi-synthetic derivate of tylosin
Product name	Tylogran® 1000 mg/g	Aivlosin® 625 mg/g	Tildosin® 250 mg/ml
Pharmaceutical form	Granulate for oral use	Granulate of use in drinking water	Solution for oral use
Presentation	Jar 550 g, jar 1 kg	Sachet 160 gram	Bottle 960 ml, can 5040 ml
Indications*	<i>Mycoplasma hyopneumoniae</i> <i>Mycoplasma hyorhinis</i>	<i>Mycoplasma hyopneumoniae</i>	<i>Mycoplasma hyopneumoniae</i>
	<i>Lawsonia intracellularis</i>	<i>Lawsonia intracellularis</i>	<i>Pasteurella multocida</i> <i>Actinobacillus pleuropneumoniae</i>
Dosage*	Lung 20 mg tylosin/kg body weight (= 2.2 g product/100 kg BW) per day, for 10 days	10 mg tylvalosin/kg body weight (= 1.6 g product/100 kg BW) per day, for 5 days	15-20 mg tilmicosine/kg body weight (= 6-8 ml product/100 kg BW) per day, for 5 days
	Intestine 5-10 mg tylosin/kg body weight (= 0.55-1.1 g product/100 kg BW) per day, for 7 days	5 mg tylvalosin/kg body weight (= 0.8 g product/100 kg LG) per day, for 5 days	n/a
Withdrawal period*	1 day	2 days	14 days

* all product information is based on the European marketing authorisations.

Read the SPC or label/leaflet before use. For more information, please contact your account manager.

Dopharma International B.V., Zalmweg 24, 4941 VX Raamsdonksveer, the Netherlands, T +31 (0)162-582000, international@dopharma.com