



Susceptibility of porcine *Streptococcus suis* isolates: MIC-values for penicillin V and penicillin G

Introduction

According to Good Veterinary Practice and the 'One Health'-concept, narrow-spectrum antimicrobial treatment is preferred above broad-spectrum treatment. Phenoxyphen® (phenoxymethylpenicillin or penicillin V) is the first narrow-spectrum penicillin for use in drinking water of pigs available in the EU. To determine the susceptibility of *S. suis* strains towards narrow-spectrum penicillins, the Minimal Inhibitory Concentrations (MIC) of *S. suis* isolates from diseased pigs were investigated. The MIC is the lowest concentration of an antimicrobial substance that, under defined *in vitro* conditions, prevents the growth of a specific bacterial strain within a defined time period.

Materials & Methods

A total of 37 unrelated *S. suis* isolates from diseased pigs were obtained from different veterinary diagnostic laboratories. The isolates were identified by MALDI-TOF MS and PCR-tests. MIC-values were determined using broth microdilution methods according to CLSI-standards^{3,4,7,8}.

14 isolates from diseased pigs in Belgium were obtained between 2010 and 2014¹. For these isolates the MIC for penicillin V was determined (0.06 µg/ml to 128 µg/ml)². 23 isolates from diseased pigs in Belgium, Germany, The Netherlands and Romania were obtained between 2019 and 2024⁵. For these isolates the MIC for penicillin G was determined (0.0625 µg/ml to 4 µg/ml)⁶.

Results

The results are summarized in table 1.

Antibiotic	<i>S. suis</i> from 2010 - 2014 (n=37)														MIC ₅₀	MIC ₉₀	S / I / R ⁷
	≤0.0625	0.125	0.25	0.5	1	2	4	8	16	32	64	128					
Penicillin V/G	31	3	3	3	0	0	0/-	0/-	0/-	0/-	0/-	0/-	0/-	≤0.0625	0.125	92% / 8% / 0%	

Table 1: MIC-distribution (penicillin G or V) of 37 *S. suis* isolates from diseased pigs in Belgium, Germany, The Netherlands and Romania, MIC50, MIC90 and the percentage of susceptible (S), intermediate (I) and resistant (R) strains.

92% (34 out of 37 isolates) of the *S. suis* strains showed a MIC of ≤ 0.125 µg/ml and were considered fully susceptible to penicillin treatment. 8% (3 out of 37 isolates) of the strains showed an intermediate susceptibility of 0.5 µg/ml. None of the investigated strains showed resistance to penicillins (MIC of ≥ 1 µg/ml).

Conclusions & Discussion

To determine the susceptibility of *S. suis* strains towards narrow-spectrum penicillins, such as Phenoxyphen[®], the Minimal Inhibitory Concentrations (MIC) of *S. suis* isolates from diseased pigs were investigated. According to the CLSI-breakpoints⁷, strains with a MIC of ≤ 0.25 $\mu\text{g/ml}$ are considered to belong to the wild type population and fully susceptible to penicillin treatment. Strains with a MIC of ≥ 1 $\mu\text{g/ml}$ are considered to have acquired genetic alterations, which result in resistance to penicillin treatment. Strains with a MIC of 0.5 $\mu\text{g/ml}$ are considered to be intermediately susceptible. Infections due to isolates with intermediate susceptibility may still be treatable with penicillin.

The 37 strains from the four European countries were highly susceptible to penicillin G or penicillin V. This investigation concerned only a limited number of *S. suis* strains. However, these results correspond to other published results as shown in table 2 and 3.

Antibiotic	<i>S. suis</i> from 2015 - 2016 (n=131)														
	0.0156	0.0313	0.0625	0.125	0.25	0.5	1	2	4	8	16	32	MIC ₅₀	MIC ₉₀	S / I / R ⁷
Penicillin G	12	66	29	5	5	6	3	1	1	1	0	2	0.0313	0.5	89% / 5% / 6%

Table 2: MIC-distribution of 131 *S. suis* isolates from diseased pigs from 5-9 European countries⁹, MIC50, MIC90 and the percentage of susceptible (S), intermediate (I) and resistant (R) strains.

Antibiotic	<i>S. suis</i> from 2024 (n=473)															
	0.0156	0.0313	0.0625	0.125	0.25	0.5	1	2	4	8	16	32	64	MIC ₅₀	MIC ₉₀	S / I / R ⁷
Penicillin G	50	258	101	30	12	13	3	4	2	0	0	0	0	0.0313	0.125	96% / 3% / 2%

Table 3: MIC-distribution of 473 *S. suis* isolates from diseased pig from The Netherlands¹⁰, MIC50, MIC90 and the percentage of susceptible (S), intermediate (I) and resistant (R) strains.

While sensitivity to penicillin G and penicillin V is virtually identical, the suitability for administration via drinking water differs for both natural penicillins. Unlike penicillin G, penicillin V is highly water-soluble and acid stable. As a result, penicillin V retains its activity after passing through the acidic environment of the stomach. Together with the high susceptibility of streptococci to penicillins, it makes the innovative registration of Phenoxyphen[®] a valuable addition to metaphylaxis and treatment options for *S. suis* infections in swine.

References

1. Diergezondheidszorg Vlaanderen.
2. Boyen, F. (2024). *The determination of minimum inhibitory concentrations of phenoxymethylpenicillin on porcine Streptococcus suis isolates*. Faculty of Veterinary Medicine, Ghent University. Registration dossier Phenoxyphen WSP.
3. Clinical and Laboratory Standards Institute (CLSI). (2013). *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals*. 4th Edition. CLSI Vet01-A4.
4. Clinical and Laboratory Standards Institute (CLSI). (2013). *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals; second informational supplement*. CLSI (Volume 28, No. 8) Vet01-S1, 2013. Editors: Watts et al. ISBN: 1-56238-880-0.
5. RIPAC-LABOR GmbH.
6. Royal GD. (2024). *Antimicrobial susceptibility testing using broth microdilution according to CLSI document VET01S*. Registration dossier Phenoxyphen WSP
7. Clinical and Laboratory Standards Institute (CLSI). (2020). *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals*. 5th Edition. CLSI supplement VET01S. Clinical and Laboratory Standards Institute, UUSA.
8. Clinical and Laboratory Standards Institute (CLSI). (2023). *Performance Standards for Antimicrobial Disk and Dilution Susceptibility Tests for Bacteria Isolated From Animals*. 6th Edition. CLSI supplement VET01S. Clinical and Laboratory Standards Institute, UUSA.
9. De Jong, A., et al. (2023). *Antimicrobial susceptibility among respiratory tract pathogens isolated from diseased cattle and pigs from different parts of Europe*. Journal of Applied Microbiology. 134, 1-14. <https://doi.org/10.1093/jambio/txad132>
10. Royal GD. (2025). *Monitoring Animal Health*.

